

## [Mini Review]

## Need-to-know basics about Zika virus infection for dental professionals

Puja NEOPANE<sup>1</sup>, Durga PAUDEL<sup>1</sup>, Koki YOSHIDA<sup>1</sup>, Bhoj Raj ADHIKARI<sup>1</sup>, Tetsuro MORIKAWA<sup>1</sup>, Aya ONISHI<sup>1</sup>,  
Daichi HIRAKI<sup>1</sup>, Osamu UEHARA<sup>2</sup>, Jun SATO<sup>1</sup>, Michiko NISHIMURA<sup>1</sup>, Itsuo CHIBA<sup>2</sup>, Yoshihiro ABIKO<sup>1\*</sup>

<sup>1</sup> Division of Oral Medicine and Pathology, Department of Human Biology and Pathophysiology,  
School of Dentistry, Health Sciences University of Hokkaido

<sup>2</sup> Division of Disease Control and Molecular Epidemiology, Department of Oral Growth and Development,  
School of Dentistry, Health Sciences University of Hokkaido

**Key words** : Zika virus infection, dental professionals, saliva,

## Abstract

Zika virus (ZIKV) infection, which spreads through the bite of an infected mosquito (*Aedes* species), is a global public health problem. Although about 80% of patients remain asymptomatic, infected individuals demonstrate common features such as fever, cutaneous rash, arthralgia, and conjunctivitis. ZIKV is known to be associated with neurological complications, such as microcephaly and Guillain-Barre syndrome. The virus spreads through contact with blood, semen, and vaginal fluids. However, it has been detected in many other human body fluids, including urine, amniotic fluid, breast

milk, saliva, cerebrospinal fluid, and cervical mucous. As dental professionals frequently come in contact with saliva and blood, it is imperative that they protect themselves from ZIKV-infected patients and instruct them to avoid the transmission of this virus. In addition, saliva can be used as a diagnostic tool for ZIKV infection. A saliva-based test may prove effective, and dental professionals might be expected to play a key role in the future surveillance and detection of the virus.

## Introduction

Zika virus (ZIKV) infection is an emerging pandemic viral disease that has been declared as a public health emergency by the World Health Organization (WHO, 2016). It has raised public concern from all around the globe due to the widespread nature of the infection at an alarming pace.

ZIKV infection is a mosquito-borne disease, which spreads primarily through the bite of an infected mosquito (*Aedes* species) (Fauci & Morens, 2016). The other routes of spread include intrauterine transmission, sexual contact, and blood transfusion (Besnard et al., 2014 ; D'Ortenzio et al., 2016 ; Musso et al, 2014). Although the majority of patients show mild symptoms such as fever, conjunctivitis, cutaneous rash, and arthralgia, this infection often causes birth defects, growth and developmental anomalies, including microcephaly, and some motor neurological manifestations in adults (Cao-Lormeau et al., 2016). Due to the increasing

number of cases of microcephaly associated with ZIKV, it can significantly affect the dental and general health of the patient. Since the ZIKV might be transmitted through body fluids including blood and saliva (Musso et al, 2014 ; Musso et al., 2015), dental practitioners should be made aware of the risk of infection. Saliva obtained from patients may be useful for a diagnostic approach. Reports about the involvement of ZIKV infection in dentistry are limited. Therefore, this review aimed to demonstrate and discuss the importance of awareness about ZIKV infection in the dental setting.

## Epidemiology

ZIKV originated from the Zika forest in Uganda and was isolated from a rhesus monkey in 1947 (Dick, 1953). The first human case of ZIKV was detected in 1952, after which it spread to other geographical regions such as Egypt (Smithburn KC et al., 1954), East Africa (Smithburn KC, 1952), Nigeria (Macnamara FN, 1954), India (Smithburn KC

et al, 1954), Japan (Taira M et al., 2017), Thailand (Pond WL, 1963), Vietnam (Smithburn KC et al, 1954), the Philippines (Hammon et al., 1958) and Malaysia (Smithburn KC et al, 1954), affecting more than 80 countries worldwide. During the outbreak of ZIKV in 2007 on several islands in the state of Yap, 5000 people were affected, while the outbreak in French Polynesia in 2013–2014 resulted in a total of 32,000 people who were affected by the virus (Schuler–Faccini et al., 2016). Thousands of babies were born with devastating birth defects after their mothers were infected in pregnancy. In 2015, a sudden increase in the cases of microcephaly in infants and the association of the Guillain–Barre syndrome was reported in Brazil (Schuler–Faccini et al., 2016). By 2016, almost 4300 cases of microcephaly had been recorded (Victora et al., 2016). However, by 2017, the number of infections had decreased owing to the several precautions that were taken into consideration. The WHO counted 12 outbreaks worldwide in 2015, 22 in 2016, and just one in early 2017. The Centers for Disease Control (CDC) and Prevention in the United States of America (US) recently deactivated its emergency response system for ZIKV that was launched in January 2016 (WHO, 2017). Although the number of infected cases has decreased, it is not completely controlled and can result in another outbreak in future. A revised guideline has been adapted by the European Centre for Disease Prevention and Control (ECDC) and WHO as a country–wise classification (WHO, 2017).

### Mode of transmission of ZIKV

ZIKV is mainly transmitted by the bite of the *Aedes* mosquito (Marchette et al., 1969). Transplacental transmission or transmission during delivery by an infected mother has also been reported (Besnard et al., 2014). Zika viral RNA has been detected in breast milk, which raises the concern of potential transmission by breastfeeding (Besnard et al., 2014). Sexual transmission was reported in a few instances (Musso et al., 2015 ; Fig.1). The Zika viral RNA has also been detected in urine at higher loads, which may hint at possibility of a person–to–person transmission route ; in addition, it indicates the possibility of testing the urine after the early symptoms of viraemia have subsided (Fauci & Morens, 2016). The virus has been found to be transmitted via blood transfusion (Musso et al., 2014). A recent study indicated that the viral RNA was detected in blood samples from asymptomatic donors, and the number of positive results was


unexpectedly high (3% ; Musso et al., 2014). On February 16, 2016, the Food and Drug Administration in the US recommended the deferral of blood donations from individuals who had traveled to areas with active ZIKV transmission, were potentially exposed to the virus, or had experienced a confirmed ZIKV infection (Zuanazzi et al., 2017).

### Signs, symptoms, and complications

ZIKV is characterized as an asymptomatic or mild dengue–like disease with fever, headache, myalgia, muscle and joint aches, conjunctivitis, and maculopapular rashes, with an incubation period of 3 to 12 days (Zuanazzi et al., 2017 ; CDC, 2016 ; Fig.1). Only about 20% of cases are symptomatic (CDC, 2016). There is no specific anti–viral treatment for this condition, and the management of symptoms may include the use of analgesics and antipyretics. However, over the period of 60 years of observation, ZIKV has not been reported to cause hemorrhagic fever or death (Fauci & Morens, 2016). The most significant public health concern is the recently confirmed association between ZIKV infection during pregnancy and microcephaly or other severe fetal brain defects (Schuler–Faccini et al., 2016). The relationship between ZIKV infection and microcephaly has long been suspected, based on the detection of Zika viral RNA in mothers and in amniotic fluid samples from fetuses, which indicates the potential of this virus to infect fetuses. Incomplete viral autophagy, centrosome abnormalities, and chromosomal instability have been observed in Zika–infected cells. Notably, no other flavivirus has been associated with teratogenic effects in humans. Guillain–Barré syndrome, an autoimmune disease causing acute or subacute flaccid paralysis, has also been found to be associated with ZIKV infection (Cao–Lormeau et al., 2016 ; Zuanazzi D et al., 2017 ; CDC, 2016).


### Detection of ZIKV in saliva

Although the transmission of ZIKV through saliva has not been substantiated so far, its presence in saliva has been detected (Musso et al., 2015). Exposure to the saliva of infected patients can be considered as a possible, yet unlikely, route of infection. The virus has been recovered in cell cultures from the saliva of infected patients and has also been detected in saliva from patients with Zika fever, especially during the acute phase of the disease (Musso et al., 2015 ; Bonaldo et al., 2016). ZIKV peptides can be detected in the



**Figure 1.** Mode of transmission of Zika virus

a. Vector borne transmission. Zika virus is transmitted by the bite of the *Aedes* mosquito. When a healthy mosquito bites a Zika infected patient/monkey, it sucks the virus-infected blood. The virus replicates inside mosquito gut, and is then transmitted to a healthy human when it is bitten by the carrier mosquito.



**Non vector-borne**

b. Non-vector born transmission. Human-to-human transmission through sexual transmission, perinatal and blood transfusion. Monkey to human transmission through the bite of a Zika-infected monkey. The clinical features include headache/fever, conjunctivitis, and muscle/joint pain. Complications such as microcephaly in new-born babies and Guillain-Barre syndrome in adults may occur.

saliva of patients previously diagnosed with Zika fever, indicating that ZIKV may be latent in the salivary gland (Balthesen et al., 1994). Viruses such as herpes simplex virus types 1 and 2 (HSV1, HSV2) and cytomegalovirus (CMV) can be transmitted through saliva (Barzon et al., 2016). Saliva may play a critical role in the human-to-human transmission of ZIKV. Nevertheless, to the best of our knowledge, there is no information about the transmission of Zika via saliva in the literature. Thus, further studies are needed to determine this phenomenon.

From another point of view, saliva can be used as a diagnostic sample for ZIKV infection (Bonardo et al., 2016). In fact, it has been already used as a source for ZIKV diagnostics when the first cases were reported during the 2014 outbreak in French Polynesia (Barzon et al., 2016). Recently, many reports confirmed that viral load (as determined by RNA levels) is significantly higher in saliva than in plasma; ZIKV RNA could be detected for a much longer period in saliva than in plasma (Barzon et al., 2016). Moreover, it was detected in the saliva of the patient even during the convalescent phase (Balthesen et al., 1994). A saliva-based test may be effective, and dental professionals may play a key role in the future surveillance and detection of this virus (Balthesen et al., 1994).

### Oral manifestations in patients with ZIKV infection

Several dental problems such as delay in the eruption of deciduous dentition, hypersalivation, higher chances of caries development, and the presence of enamel alterations consistent with dental hypoplasia have been observed in children with microcephaly associated with prenatal ZIKV infection (Cavalcanti, 2017). Therefore, it is essential for the dental professional to provide proper attention to the patient during dental care and adequate information regarding oral hygiene and eating habits to the caretakers and family. Severe neurological complications associated with Guillain-Barré syndrome have been described in postnatal ZIKV infections (Passi et al., 2017). Patients with Guillain-Barré syndrome present with muscle weakness caused due to the damage of the peripheral nervous system by the immune system. Facial muscle weakness with disorders of muscle movement and swallowing are often observed in these patients (Rajabally et al., 2012). Nonetheless, no specific orofacial manifestations have been reported so far. A recent pa-

per reported the presence of hyperemia and petichiae in the hard palate of a patient with ZIKV. Since ZIKV are neurotropic viruses, which are capable of infecting and replicating in neural cells (Brasil et al., 2016), they may persist in the latent state and are reactivated at a later period, as in the cases of HSV that cause recurrent ulcerations in the oral mucosa. ZIKV could invade endothelial cells, fibroblasts, and keratinocytes via receptors in the oral mucosal tissues (Brasil et al., 2016), causing specific oral pathogenic conditions. However, further investigations are needed to determine the presence of oral specific manifestations.

### Role of dental professionals for ZIKV infection control

No appropriate treatment strategy or vaccine for ZIKV is available thus far. Protection of dentists from the infection and avoiding the transmission of this condition to other patients is crucial. Both saliva and blood are known to be the routes for transmission of ZIKV infection during dental procedures. Active participation of dental professionals for the screening of ZIKV infected patients and their adherence to infection control guidelines are of utmost importance in controlling the dissemination of ZIKV infection. It is advisable for dental professionals, who encounter suspected patients with ZIKV infections, to report them to the local health department for further investigations (CDC, 2016).

### References

- Balthesen M, Dreher L, Lucin P, Reddehase MJ. The establishment of cytomegalovirus latency in organs is not linked to local virus production during primary infection. *J Gen Virol* 75 : 2329–2336, 1994.
- Barzon L, Pacenti M, Berto A, Sinigaglia A, Franchin E, Lavezzo E, Brugnaro P, Palu G. Isolation of infectious Zika virus from saliva and prolonged viral RNA shedding in a traveller returning from the Dominican Republic to Italy, January 2016. *Euro Surveill* 21 : 30159, 2016.
- Besnard M, Lastère S, Teissier A, Cao-Lormeau V, Musso D. Evidence of perinatal transmission of Zika virus, French Polynesia, December 2013 and February 2014. *Euro Surveill* 19 : 20751, 2014.
- Bonardo MC, Ribeiro IP, Lima NS, Dos Santos AA, Meneses LS, da Cruz SO, de Mello IS, Furtado ND, de Moura EE, Damasceno L, da Silva KA, de Castro MG, Gerber AL, de Almeida LG, Lourenço-de-Oliveira R,

- Vasconcelos AT, Brasil P. Isolation of Infective Zika Virus from Urine and Saliva of Patients in Brazil. *PLoS Negl Trop Dis* 10 : e0004816, 2016.
- Brasil P, Calvet GA, de Souza RV, Siqueira AM. Exanthema associated with Zika virus infection. *Lancet Infect Dis* 16 : 866, 2016
- Cao-Lormeau VM, Blake A, Mons S, Lastère S, Roche C, Vanhomwegen J, Dub T, Baudouin L, Teissier A, Larre P, Vial AL, Decam C, Choumet V, Halstead SK, Willison HJ, Musset L, Manuguerra JC, Despres P, Fournier E, Mallet HP, Musso D, Fontanet A, Neil J, Ghawché F. Guillain-Barré syndrome outbreak associated with Zika virus infection in French Polynesia : a case-control study. *Lancet* 387 : 1531-1539, 2016.
- Cavalcanti AL. Challenges of Dental Care for Children with Microcephaly Carrying Zika Congenital Syndrome. *Contemp Clin Dent* 8 : 345-346, 2017.
- Centers for Disease Control and Prevention. Zika Virus : For health care providers : Clinical Evaluation and Disease. Available from : [http://www.cdc.gov/zika/hc-providers/clinical\\_evaluation.html](http://www.cdc.gov/zika/hc-providers/clinical_evaluation.html)(accessed on April 13, 2016).
- Council of State and Territorial Epidemiologists. Notice to Readers : Changes in the Presentation of Zika Virus Disease, Non-Congenital Infection, and Addition of Zika Virus Congenital Infection to Notifiable Diseases and Mortality Table I. Available at : [cdc.gov/mmwr/volumes/65/wr/mm6520a6.htm](http://cdc.gov/mmwr/volumes/65/wr/mm6520a6.htm). 65 : 524, 2016.
- D'Ortenzio E, Matheron S, Yazdanpanah Y, de Lamballerie X, de Lamballerie X, Hubert B, Piorkowski G, Maquart M, Descamps D, Damond F, Leparç-Goffart I. Evidence of sexual transmission of Zika virus. *N Engl J Med* 374 : 2195-2198, 2016.
- Dick GW. Epidemiological notes on some viruses isolated in Uganda ; Yellow fever, Rift Valley fever, Bwamba fever, West Nile, Mengo, Semliki forest, Bunyamwera, Ntaya, Uganda S and Zika viruses. *Trans R Soc Trop Med Hyg* 47 : 13-48, 1953.
- Fauci AS & Morens DM. Zika virus in the Americas — yet another arbovirus threat. *N Engl J Med* 377 : 601-604, 2016.
- Hammon WM, Schrack WD Jr. & Sather GE. Serological survey for a arthropod-borne virus infections in the Philippines. *Am J Trop Med Hyg* 7 : 323-328, 1958.
- Macnamara FN. Zika virus : A report on three cases of human infection during an epidemic of jaundice in Nigeria. *Trans R Soc Trop Med Hyg* 48 : 139-145, 1954.
- Marchette NJ, Garcia R & Rudnick A. Isolation of Zika virus from *Aedes aegypti* mosquitoes in Malaysia. *Am J Trop Med Hyg* 18 b : 411-415, 1969.
- Musso D, Nhan T, Robin E, Roche C, Bierlaire D, Zisou K, Shan Yan A, Cao-Lormeau VM, Broult J. Potential for Zika virus transmission through blood transfusion demonstrated during an outbreak in French Polynesia, November 2013 to February 2014. *Euro Surveill* 19 : 1-3, 2014.
- Musso D, Roche C, Nhan TX, Robin E, Teissier A, Cao-Lormeau VM. Detection of Zika virus in saliva. *J Clin Virol* 68 : 53-55, 2015.
- Musso D, Roche C, Robin E, Nhan T, Teissier A, Cao-Lormeau VM. Potential sexual transmission of Zika virus. *Emerg Infect Dis* 21 : 359-361, 2015.
- Passi D, Sharma S, Dutta SR, Ahmed M. Zika virus diseases – The new face of an ancient enemy as global public health emergency (2016) : Brief review and recent updates. *Int J Prev Med* 8 : 6, 2017.
- Pond WL. Arthropod-borne virus antibodies in sera from residents of South-East Asia. *Trans R Soc Trop Med Hyg* 57 : 364-371, 1963.
- Rajabally, Y.A. & Uncini, A. Outcome and its predictors in Guillain-Barr syndrome. *J Neurol Neurosurg Psychiatry*, 83 : 711-718, 2012.
- Schuler-Faccini L, Ribeiro EM, Feitosa IM, Horovitz DD, Cavalcanti DP, Pessoa A, Doriqui MJ, Neri JI, Neto JM, Wanderley HY, Cernach M, El-Husny AS, Pone MV, Seroa CL, Sanseverino MT. Possible association between Zika virus infection and microcephaly – Brazil, 2015. *MMWR Morb Mortal Wkly Rep* 65 : 59-62, 2016.
- Smithburn KC. Neutralizing antibodies against arthropod-borne viruses in the sera of long-time residents of Malaya and Borneo. *Am J Hyg* 59 : 157-163, 1954.
- Smithburn KC, Taylor RM, Rizk F & Kader A. Immunity to certain arthropod-borne viruses among indigenous residents of Egypt. *Am J Trop Med Hyg* 3 : 9-18, 1954.
- Smithburn KC. Neutralizing antibodies against certain recently isolated viruses in the sera of human beings residing in East Africa. *J Immunol* 69 : 223-234, 1952.
- Smithburn KC, Kerr JA & Gatne PB. Neutralizing antibodies against certain viruses in the sera of residents of India. *J Immunol* 72 : 248-257, 1954.

Taira M, Ogawa T, Nishijima H, Yamamoto K, Hotta C, Akita M, Tajima S, Saijo M. The First Case of Zika Virus Isolated from a Japanese Patient Who Returned to Japan from Fiji in 2016. *Jpn J Infect Dis* 70 : 586–589, 2017.

Victora CG, Schuler-Faccini L, Matijasevich A, Ribeiro E, Pessoa A, Barros FC. Microcephaly in Brazil : How to interpret reported numbers? *Lancet* 387 : 621–624, 2016.

World Health Organization [WHO] (2017b). *Zika Virus Situation Reports*. Available at : <http://www.who.int/emergencies/zika-virus/situation-report/en/> World Health Organization Zika virus country classification scheme, March 2017.

Zika virus tied to birth defects is international emergency, WHO says. Yahoo Sports. [http://sports.yahoo.com/news/declares-zika-virus-global-public-health-emergency-184358252finance.html;\\_ylt=AwrTceNhra9Wu\\_oAouwnnllQ;\\_ylu=X3oDMTByNWU4cGh1BGNvbG8DZ3ExBHBvcwMxBHZ0aWQDBHNIYwNzYw--](http://sports.yahoo.com/news/declares-zika-virus-global-public-health-emergency-184358252finance.html;_ylt=AwrTceNhra9Wu_oAouwnnllQ;_ylu=X3oDMTByNWU4cGh1BGNvbG8DZ3ExBHBvcwMxBHZ0aWQDBHNIYwNzYw--). Published February 1, 2016.

Zuanazzi D, Arts EJ, Jorge PK, Mulyar Y, Gibson R, Xiao Y, Bringel Dos Santos M, Machado MAAM, Siqueira WL. Postnatal Identification of Zika Virus Peptides from Saliva. *J Dent Res* 96 : 1078–1084, 2017.



Puja NEOPANE

**Education :**

2005 : Graduated from Rajiv Gandhi University of Health Sciences, Bangalore, India (Bachelor of Medical Laboratory Technology)

2015 : Post graduation from Tribhuvan University, Nepal (MSc. Clinical Microbiology)

2016 : Graduate student in Division of Oral medicine and Pathology, Health Sciences University of Hokkaido, Japan. (running)

**Professional Experiences :**

2006 to 2012 : Senior medical laboratory technologist, Medicare National Hospital and Research Centre Ltd, Kathmandu, Nepal.

2015 : Teaching assistant for medical student, Chitwan Medical College and Teaching Hospital, Nepal.