

[REVIEW]

Effectiveness of Bee Pollen as an Anti-Aging Agent

Ariuntsetseg KHURELCHULUUN^{1,2}, Durga PAUDEL³, Osamu UEHARA⁴, Takahashi SHUHEI¹, Dedy ARIWANSA¹, Koki YOSHIDA¹, Tetsuro MORIKAWA¹, Fumiya HARADA², Jun SATO¹, Hiroko MIURA⁴, Hiroki NAGAYASU², Yoshihiro ABIKO¹

¹Division of Oral Medicine and Pathology, Department of Human Biology and Pathophysiology, School of Dentistry, Health Sciences University of Hokkaido

²Division of Oral and Maxillofacial Surgery, Department of Human Biology and Pathophysiology, School of Dentistry, Health Sciences University of Hokkaido

³Advanced Research Promotion Center, Health Sciences University of Hokkaido

⁴Division of Disease Control and Molecular Epidemiology, Department of Oral Growth and Development, School of Dentistry, Health Sciences University of Hokkaido

Key words : bee pollen, anti-aging, bioactive compound

Abstract

Bee pollen has been used as a food and supplement for a long time. Bee pollen in diet can bring many beneficial effects on human health. Aging is an inevitable progress which leads to age-related degenerative diseases. Most of the age-related diseases are attributed to the accumulated oxidative stress in tissue over years.

Natural products such as bee pollen are rich in antioxidant and anti-inflammatory activity contributing to anti-aging effect. Bee pollen contains abundant phenolic compounds, which are highly efficient antioxidants. Herein, we summarize the potential use of bee pollen as an anti-aging component.

Introduction

A variety of bee products have been used as food and medicine since ancient days. Products such as honey, propolis, bee pollen, royal jelly, beeswax, and venom have been studied for their beneficial effects on human health (Pasupuleti et al., 2017). Bee pollen abundant in proteins, amino acids, phenolic acids, and flavonoids (Campos et al., 2008), which have antioxidant, anti-inflammatory, antibacterial, and anticancer activities.

Aging is a complex process in humans and other living organisms. It is associated with physiological and functional decline, which leads to the development of age-related diseases such as arterial sclerosis, diabetes, cataract, Parkinson's disease, Alzheimer's disease, and others (Jaul and Barron, 2017). The acceleration of the aging process due to individual lifestyles may promote the development of age-related diseases (Rescigno et al., 2017). The accumulation of reactive oxygen species over a period of time is one of the most important factors for the acceleration of the aging process (Oliveira et al., 2010). The scavenging of the accu-

mulated reactive oxygen species, which is an antioxidant activity, may serve as an anti-aging process (Wojdyło et al., 2018). Hence, owing to its antioxidant properties, bee pollen is proved to be beneficial as a potential natural anti-aging supplement.

Although a few studies have demonstrated the anti-aging effect of bee pollen *in vivo* and *in vitro*, a comprehensive review is lacking (Graikou et al., 2008 ; Liu and Li, 1990). The antioxidant property of bee pollen is due to the presence of its bioactive components, such as phenolics and flavonoids. The concentrations of these components in bee pollen vary among geographical regions (Kaškonienė et al., 2015).

The aim of this review is to summarize the potential use of bee pollen as an anti-aging component. Additionally, variations in the amounts of phenolics and flavonoids and the antioxidant properties of bee pollen are discussed.

Components of bee pollen

Bee pollen is produced by the stamen of the flower and collected by bees. The structure and ratio of the bioactive

components in pollen differ among the various types of plants (Leja et al., 2007). The pollen grain consists of two layers, an outer shell, and the inner cytoplasm, which house a tube cell and a generative cell. The contents of the bioactive components in pollen can be affected by climate conditions, seasonal changes, or the use of pesticides.

Bees are the most important pollinators for all plants/flowers (Hellerstein et al., 2017). During foraging, the pollen sticks onto the hairy bodies of the worker bees and is transferred to a pollen basket on the hind limbs where it is agglutinated with bee enzymes to form bee pollen. The bee pollen is carried to beehives where they are stored in niches of honeycombs as the main food for the bee colonies. Thus, the bee pollen should be collected before the worker bees enter the hive.

Various bee pollen products are available over-the-counter as supplements or food additives, and they are broadly divided into three major forms (Nuvoloni et al., 2020). The first is the raw bee pollen granules that are directly collected from the bee hive (Nogueira et al., 2012). The second type comprises granules that are milled down to a powdered form, which can easily dissolve in liquids. The third type is the bee pollen extract, which has the highest nutritional value. The color of bee pollen varies from bright yellow to dark green or brown depending on its composition.

To date, over 200 chemical substances have been found in bee pollen. The main components include proteins and amino acids, reducing sugars, lipids, nucleic acids, and fiber (Campos et al., 2008 ; Almeida–Muradian et al., 2005). The minerals present include potassium, sodium, calcium, iron, zinc, and copper, and the vitamins include provitamin A (β -carotene), vitamin E (tocopherol), niacin, thiamine, biotin, folic acid, and enzymes. Bioactive substrates, such as phenolic compounds (phenolics, flavonoids, anthocyanins, phenylpropanoids), fatty acids (linoleic, γ -linoleic, and arachidonic acid), phospholipids, and phytosterols (Leja et al., 2007 ; Tomás–Lorente et al., 1992) contribute to the health effects of bee pollen.

Biological properties of bee pollen

Bee pollen has many biological activities, such as antioxidant, anti-inflammatory, anticancer, and antimicrobial activities. The antioxidant activities may be directly involved in the anti-aging process. The antioxidant activities of the

components in bee pollen are shown in the section “Studies on the antioxidant activity of bee pollen”.

Bee pollen possesses anti-inflammatory effects similar to other apitherapeutic bee products such as propolis, bee venom, honey, and royal jelly (Kolayli et al., 2016). Phenolic compounds in bee pollen may exert anti-inflammatory activities by modulating the B cells, mast cells, macrophages, T cells, and neutrophils (Middleton & Kandaswami, 1992).

Important inflammatory cytokines, such as TNF- α , IL-1 β , and IL-6, might be inhibited by natural phenolic compounds via macrophage cell response (González et al., 2011 ; Yahfoufi et al., 2018). Quercetin is a phenolic component commonly found in bee pollen; it inhibits the activity of cyclooxygenase and lipoxygenase during the synthesis of prostaglandins from arachidonic acid. The inhibitory effect can halt the proinflammatory process and migration of immune cells, and reduce edema (Middleton E, 1998 ; M. Choi, 2007). Nonsteroidal anti-inflammatory drugs are known to have various side effects; hence, bee pollen may be considered as an alternative anti-inflammatory medication (Bindu et al., 2020).

Studies have shown that bee pollen can act against a variety of cancer cells. Quercetin, commonly found in bee pollen, demonstrated a significant inhibitory effect on the growth of ovarian cancer cells (Vafadar., et al 2020). Another pollen extract called cernitin was found to be effective for the treatment of benign prostatic hyperplasia (Yasumoto et al., 1995). The synergistic anti-tumor activities of bee pollen have been proven in breast cancer cells; bee pollen promoted cancer cell apoptosis and the inhibition of cancer cell growth in combination with cisplatin, suggesting that it might contribute to chemotherapy (Omar et al., 2016 ; Amalia et al., 2020). The activity of caspase-3, which is the main enzyme related to the induction of apoptosis, in HL-60/promyeloblast cells was increased following stimulation with bee pollen alone (Uçar et al., 2016). Furthermore, bee pollen alone might function as a potential anti-cancer agent. However, additional studies are required to determine the clinical potential of bee pollen as an anti-cancer agent.

From ancient times, bee products have been used as a natural remedy for bacterial infections (Kurek–Górecka et al., 2020). The antimicrobial property of bee pollen might be related to the amylase enzymes produced by the salivary glands of bee (Campos et al., 2008). Amylases, which hy-

drolyze the polysaccharide backbone of extracellular polymeric substances, might contribute to the antimicrobial properties of bee pollen (Lahiri et al., 2021). Phenolic compounds such as quercetin, epicatechin, and tannic acid found in bee pollen inhibit β -lactamase activity and the formation of the bacterial biofilm (Mandal et al., 2017). Moreover, the phenolic compounds in bee pollen may be involved in antimicrobial activities. In our previous study, mice fed with a 5% bee pollen diet showed an increase in the abundance of beneficial bacteria such as *Lactobacillus* in the oral cavity and intestine; in addition, the growth of the periodontal pathogen *Porphyromonas gingivalis* was inhibited at a concentration of >2.5% (Khurelchuluun et al., 2021). Another study showed the diverse effects of bee pollen on various microorganisms, with *Candida albicans* being the most resistant and *Staphylococcus epidermidis* being the most sensitive (Soares de Arruda et al., 2021). These findings indicate that the antimicrobial activity of bee pollen might be dependent on the species of bacteria.

Function of natural foods as an anti-aging agent

Natural foods rich in bioactive components are known to have anti-aging activities. Bioactive components such as phenolic compounds, monounsaturated fat, and vitamins are the main antioxidants sources (Chrysohoou & Stefanadis, 2013). Curcumin, royal jelly, and blueberry are rich in bioactive components. Curcumin/turmeric is one of the main spices used in the Middle East. Diarylheptanoid, the main active ingredient in curcumin, is a strong antioxidant consisting of two aromatic rings. In animals, curcumin was found to extend the life span, protect against oxidative stress, and modulate the expression levels of several age-related genes (Lee et al., 2010). Royal jelly is another bee product and is similar to bee pollen. The anti-aging potential of royal jelly has been studied previously; a bee/queen bee fed with royal jelly was found to live 20 times longer than other bees (De Loof, 2011). Enhancement of the antioxidative capacity was suggested to extend the lifespan in various *in vivo* animal studies (Kunugi and Mohammed Ali, 2019). Blueberry has the most abundant antioxidant compounds among fruits. Some studies reported that the lifespan-prolonging activity of blueberry might involve the regulation of the related genes and the promotion of the Phase 2 defense enzymes superoxide dismutase (SOD) and catalase (CAT) (Peng et al.,

2012; Wojdyło et al., 2018). Certain environmental factors such as stress, UV exposure, pollution, and poor diet contribute to accelerated aging. Aging significantly decreases the quality of life and raises the risk of age-related diseases, such as neurodegenerative disease, cardiovascular disease, arthritis, hypertension, and diabetes (Jaul & Barron, 2017). Natural foods related to anti-aging have been successfully used to mitigate the harmful effects of environmental factors during aging.

The anti-aging potential of many foods is explained by their high content of phenolic compounds (Luceri et al., 2017). Phenolic compounds can be classified into two broad groups: phenolics and polyphenols/flavonoids. Phenolic compounds contain aromatic rings with a hydroxyl group in their molecular structure and are known as the secondary metabolites of plants. They possess the highest antioxidant potential among all the other natural substances (Safar et al., 2015). The degree of antioxidant activity is related to the number of hydroxyl groups. The hydrogen atom from the hydroxyl group attaches to the active free radicals within the cell. Free radicals or ROS include highly active ions such as superoxide anion and hydroxyl radical, while the nonradical oxygen derivatives include singlet oxygen and stable hydrogen peroxide.

Excessive ROS can act as a secondary messenger in cellular signaling cascades and trigger prolonged DNA damage to the cell. Although ROS plays a key role in the normal biological developmental processes in cells, excessive ROS leads to the oxidative stress. Oxidative stress is a pathological condition caused due to an imbalance in the ROS and the antioxidant substances (Barnham et al., 2004).

Bee pollen is rich in phenolic and flavonoid components, which contribute to the antioxidant properties (Morais et al., 2011). These properties may be responsible for the anti-aging action of bee pollen.

Studies on the antioxidant activity of bee pollen

We summarized the bee pollen studies in Table 1. Studies are performed total phenolic content, total flavonoid content with antioxidant activity of various bee pollen. The antioxidant activity in bee pollen has been evaluated by different methods, such as the 1,1-diphenyl-2-picryl-hydrazil (DPPH) assay, ABTS-radical scavenging capacity, and beta-carotene/linoleic bleaching test (Table 1). The wide vari-

Table 1. Summary of the studies on antioxidant activity of bee pollen

Authors	Region and Country	Content of phenolic compound(s)		Antioxidant activity [DPPH][TE/g]
		Total phenolic content [GAE/g]	Total flavonoid content [QE/g or RE/g]	
AbdElsalam (2018)	Egypt	0.8 – 2.3 mg	0.1 – 0.85 mg	63 – 90 RSA%
Anjos (2019)	Spring of 2017 in Castelo Branco, Portugal	35.05 mg	6.99 mg QE/g	2.62 mg/mL
Atsalakis (2017)	Greek endemic	15.2 – 60.2 mg	6.0 – 57.6 mg	0.7 – 44.6 RSA%
Bárbara (2015)	Bahia, Brazil (João Dourado)	40 ± 13 mg	1.0 ± 0.2 mg	–
Carpes (2013)	Brazil (Parana, Santa Catarina)	20.2 – 48.8 mg	6.6 – 28.4 mg	14.9 – 94.7 RSA%
Čeksterytė (2016)	Lithuania	3.5 – 23.3 mg	–	1.1 – 1.5 mg
De-Melo (2018) a	Brazilian regions	6.5–29.2 mg	0.3–17.5 mg	9.4–155 mmol
De-Melo (2018) b	Brazilian regions (Sao Paulo, Bahia)	13.31 mg	0.3 – 19.0 mg	10.0 – 110.8 mmol
Domenici (2015)	Italy (Tuscany)	13.5 – 24.7 mg	5.91 – 15.86 mg	37.95 – 94.45 RSA%
Duarte (2018)	Alagoas, Northeast region of Brazil	6.9 – 21 mg	0.3 – 17 QE/g	76.5 – 83.3 %
Fadzilah (2017)	Malaysia	33.5 – 135.9 mg	15.3 – 31.8 mg	0.86 – 3.24 mg/mL
Fatrcová, Š (2015)	Slovakia	0.69 – 0.80 mg	0.23 – 0.61 mg	48 – 50 RSA%
Freire (2012)	Canavieiras (northeastern Brazil)	41.5 – 213.2 mg	–	2.5 – 209 mg/mL
Jin (2018)	China	–	–	IC50 : 2.29–3.48 mg/mL
Karabagias (2018)	Greece	5 mg GAE/ml	–	68.7 – 88.26 RSA%
Kaškonienė (2014)	Latvia, Lithuania, China, Spain	14.2 – 26.8 mg	2.7 – 5.2 mg	0.9 – 8.4 mg
Ketkar (2014)	West Bengal, India	18.29 mg	12.23 mg	54.79 g/mL
Kostić (2019)	Serbia	2.91 – 3.82 mg	0.84 – 0.87 mg	–
LeBlanc (2009)	North of Tucson, USA	15.91–34.85 mg	2.66–5.48 mg	–
Mohdaly (2015)	Egypt	–	–	5 ug/ml conc : 15%
Morais (2011)	North and Centre regions of Portugal	10.50–16.80 mg	–	2.24 – 5.87 mg/mL
Pascoal (2014)	Portugal, Spain	18.5 – 32.1 mg	3.9 – 10.14 mg	0.35 – 3.70 mg/mg
Rocchetti (2019)	Italy	4.2 – 29.6 mg	–	11.9 – 134.7 mol
Sun (2017)	China	12.57 mg	22.89 mg	–
Ulusoy (2014)	Turkey	~ 74 mg	–	SC50 3.3 mg/ml
Vasconcelos (2017)	Brazil	5.85–46.25 mg	1.8 – 107.0 mg	42.35 – 78.58 RSA%
Zhang (2016)	China	–	604 mg RE/g	–
Zuluaga (2016)	Colombian central region	24.8 – 33.7 mg	552 – 778 μmol	100.6 – 126.5 μmol

[GAE : gallic acid equivalent ; QE : quercetin equivalent ; RE : rutin equivalent ; DPPH : 2,2-diphenyl-1-picrylhydrazyl ; RSA : radical scavenging activity ; TE/g : μmol Trolox equivalents per gr]

ation in the antioxidant activity and the concentrations of the phenolic and flavonoid components might be attributed to geographical variations and differences in the evaluation techniques. Nonetheless, bee pollen is reported to be rich in phenolic and flavonoid components, which contribute to its high antioxidant effect (Rzepecka-Stojko et al., 2015).

Most of the studies in Tab. 1 focused on bee pollen are from Brazil and Spain, which are renowned for their bee pollen industries. Quercetin and following kaempferol, isorhamnetin, naringenin, rutin and cinnamic acid, caffeic acid, gallic acid, ferulic acid, luteolin, myricetin, catechin, and quinic acid were the most commonly found phenolic compounds in bee pollen. Quercetin, a strong antioxidant, is known to have a high effect on longevity and stress resistance in animal models (Pietsch et al., 2012).

Mechanism of antioxidant activity of bee pollen

The antioxidant activity of bee pollen involves the scavenging of excessive ROS. The phenolic compounds in bee pollen might be the main components responsible for its antioxidant activity (Leja et al., 2007). The commonly involved ROS pathways and the possible intervention by phenolic compounds in bee pollen are summarized in Figure 1.

The accumulation of excessive ROS can damage the cell via various pathways. The nuclear factor erythroid 2 (NFE2)–related factor 2 (Nrf2) is essential for signaling the production of drug-metabolizing enzymes. Nrf2 is connected to the endogenous inhibitor Kelch-like ECH associated protein 1 (KEAP1). ROS inhibits the ability of KEAP1 to attach to Nrf2, and the released Nrf2 connects to the antioxidant response element (ARE). Cytoprotective gene expression is




Figure 1. Excessive ROS pathways in oxidative stress. High amount of ROS in cells leads to cell senescence, DNA damage, cell apoptosis and cell death. Phenolic compounds in natural foods work as an antioxidant agent and halt the oxidative damages within the cell.

regulated by Nrf2–ARE binding, which codes the cellular antioxidant and anti–inflammatory cytokine defenses. Although the ROS–Nrf2 pathway is a normal biological defense system, a high amount of ROS can activate tyrosine kinases and lead to the failure of the Nrf2–Keap1 complex (Sajadimajd and Khazaei, 2018). Quercetin has been shown to enhance the Nrf2–ARE binding activity; Nrf2 protein is stabilized by the deceleration of the ubiquitination and proteasomal turnover, whereas Keap1 protein is reduced by the formation of a modified Keap1 protein in exposure of quercetin (Tanigawa et al., 2007). Phenolic compounds can induce Phase 2 enzymes such as CAT, SODs, and glutathione (GSH), these work as an endogenous antioxidant (Petri et al., 2012).

NF–κB (nuclear factor kappa–light–chain–enhancer of activated B cells) is a transcriptional factor that targets genes, particularly those involved in the antioxidant response, such as the ferritin heavy chain and SOD2. Another NF–κB target gene that contributes to both survival and innate immune functions is the HIF–1α gene, which encodes the oxygen–regulated subunit of the hypoxia–responsive transcription factor–1. Phenolic compounds can inhibit the NF–κB pathway to impart the antioxidant property (Pham et al., 2004).

Cytochrome C is a designated scavenger of ROS and is released from the mitochondrial membrane of the cell. Ex-

cessive ROS induces activation of caspase 3 and caspase–3–like proteases caused by the release of cytochrome C (Higuchi et al., 1998). The C–Jun NH (2)–terminal kinase (JNK) signaling pathway is one of the mitogen–activated protein kinase (MAPK) signaling pathways, which regulates certain cellular processes such as cell proliferation, embryonic development, and apoptosis. Over–activation of JNK/c–Jun signals induced by excessive ROS can lead to cell death (Singh et al., 2009). The inhibition of these pathways by phenolic compounds can result in antioxidative effects. P38/MAPK is the highly conserved enzyme responsive to oxidative stress. It can regulate the production of proinflammatory mediators (such as IL–1β, TNF–α, and IL–6), and is involved in cell differentiation, apoptosis, and autophagy processes. The ROS–induced p38/MAPK pathway induces the exhaustion of the cell population and limits cell proliferation (Ito et al., 2006). Phenolic compounds inhibit the p38/MAPK pathway (Xia et al., 2018). Furthermore, *p*–coumaric acid, one of the phenolic acids in bee pollen, enhances the ability of the cell to resist the SKN–1–mediated oxidative stress and regulates the activity of the MAPK pathway (Yue et al., 2019).

Matrix metalloproteinases (MMPs) are a group of enzymes responsible for the degradation of proteins and collagens in the extracellular matrix. MMPs play a major role

during all stages of the tissue healing process by modifying the structural tissues and supporting cell growth. ROS is known to trigger the MMPs via the NF- κ B pathway. Elevated MMPs are often observed in cancer cells and contribute to their growth and invasion (Morgia et al., 2005). The phenolic component, including polyphenols, phenolic acids, and monophenols act against tumor growth and invasion by reducing the MMP content (Weng and Yen, 2012).

Conclusion

Bee pollen has antioxidant properties, which can prove useful during the anti-aging process. This review demonstrates the potential use of bee pollen as a natural supplement that might aid in the prevention of pathological aging in humans.

References

- AbdElsalam E, Foda HS, Abdel-Aziz MS & Abd FK. Antioxidant and antimicrobial activities of Egyptian bee pollen. *Middle East J. Appl. Sci.* 8 : 1248–1255, 2018.
- Almeida–Muradian LB, Pamplona LC, Coimbra S & Barth OM. Chemical composition and botanical evaluation of dried bee pollen pellets. *J Food Compost Anal* 18 : 105–111, 2005.
- Amalia E, Diantini A & Subarnas A. Water soluble propolis and bee pollen of *Trigona* spp. from South Sulawesi Indonesia induce apoptosis in the human breast cancer MCF-7 cell line. *Oncol. Lett.* 20 : 1–1, 2020.
- Anjos O, Paula VB, Delgado T & Estevinho LM. Influence of the storage conditions on the quality of bee pollen. 2019.
- Atsalakis E, Chinou I, Makropoulou M, Karabournioti S & Graikou K. Evaluation of phenolic compounds in *Cistus creticus* bee pollen from Greece. Antioxidant and antimicrobial properties. *Natural Product Communications.* 12 : 11, 2017.
- Bárbara MS, Machado CS, Sodr  GD, Dias LG, Estevinho LM & De Carvalho CA. Microbiological assessment, nutritional characterization and phenolic compounds of bee pollen from *Melipona mandacaia* Smith, 1983. *Molecules* 20 : 12525–12544, 2015.
- Barnham KJ, Masters CL & Bush AI. Neurodegenerative diseases and oxidative stress. *Nat. Rev. Drug Discov.* 3 : 205–214, 2004.
- Bindu S, Mazumder S & Bandyopadhyay U. Non-steroidal anti-inflammatory drugs (NSAIDs) and organ damage : A current perspective. *Biochem Pharmacol* 180 : 114147, 2020.
- Campos MGR, Bogdanov S, de Almeida–Muradian LB, Szczesna T, Mancebo Y, Frigerio C & Ferreira F. Pollen composition and standardisation of analytical methods. *J. Apic. Res.* 47 : 154–161, 2008.
- Carpes ST, De Alencar SM, Cabral IS, Oldoni TL, Mour o GB, Haminiuk CW, Da Luz CF & Masson ML. Polyphenols and palynological origin of bee pollen of *Apis mellifera* L. from Brazil. Characterization of polyphenols of bee pollen. *CyTA–Journal of Food* 11 : 150–161, 2013.
-  eksteryt  V, Navakauskien  R, Treigytyt  G, Jansen E, Kurtinaitien  B, Dabkevi ien  G, Bal ekas J. Fatty acid profiles of monofloral clover beebread and pollen and proteomics of red clover (*Trifolium pratense*) pollen. *Bioscience, biotechnology, and biochemistry.* 80 : 2100–2108, 2016.
- Choi EM. Antinociceptive and antiinflammatory activities of pine (*Pinus densiflora*) pollen extract. *Phytother Res* 21 : 471–475, 2007.
- Chrysohoou C & Stefanadis C. Longevity and Diet. Myth or pragmatism? *Maturitas* 76 : 303–307, 2013.
- De Loof A. Longevity and aging in insects : Is reproduction costly ; cheap ; beneficial or irrelevant? A critical evaluation of the “trade-off” concept. *J Insect Physiol.* 57 : 1–11, 2011.
- De–Melo AA, Estevinho LM, Moreira MM, Delerue–Matos C, de Freitas AD, Barth OM & de Almeida–Muradian LB. A multivariate approach based on physicochemical parameters and biological potential for the botanical and geographical discrimination of Brazilian bee pollen. *Food Bioscience.* 25 : 91–110, 2018.
- De-Melo AA, Estevinho LM, Moreira MM, Delerue-Matos C, Freitas AD, Barth OM & Almeida-Muradian LB. Phenolic profile by HPLC-MS, biological potential, and nutritional value of a promising food : Monofloral bee pollen. *Journal of Food Biochemistry* 42 : e12536, 2018.
- Domenici V, Gabriele M, Parri E, Feliccioli A, Sagona S, Pozzo L, Biondi C & Pucci L. Phytochemical composition and antioxidant activity of Tuscan bee pollen of different botanic origins.
- Duarte AW, Vasconcelos MR, Oda–Souza M, Oliveira FF & L pez AM. Honey and bee pollen produced by *Meliponini* (Apidae) in Alagoas, Brazil : multivariate

- analysis of physicochemical and antioxidant profiles. *Food Science and Technology*. 11 : 493–503, 2018.
- Fatrcová–Šramková K, Nôžková J, Máriássyová M, Kačániová M. Biologically active antimicrobial and antioxidant substances in the *Helianthus annuus* L. bee pollen. *Journal of Environmental Science and Health, Part B*. 51 : 176–181, 2016.
- Freire KR, Lins A, Dórea MC, Santos FA, Camara CA & Silva T. Palynological origin, phenolic content, and antioxidant properties of honeybee–collected pollen from Bahia, Brazil. *Molecules*. 17 : 1652–1664, 2012.
- González R, Ballester I, López–Posadas R, Suárez MD, Zarzuelo A, Martínez–Augustin O & Medina FSD. Effects of Flavonoids and other Polyphenols on Inflammation. *Crit. Rev. Food Sci. Nutr*. 51 : 331–362, 2011.
- Graikou K, Aligiannis N, Chondrogianni N, Kapeta S, Gonos ES & Chinou. Chemical analysis and anti–aging properties of Greek pollen. *Planta Med* 74 : PC64, 2008.
- Harif FN, Jaapar MF, Jajuli R & Wan Omar WA. Total phenolic content, total flavonoid and antioxidant activity of ethanolic bee pollen extracts from three species of Malaysian stingless bee. *Journal of Apicultural Research*. 56 : 130–135, 2017.
- Hellerstein D, Hitaj C, Smith D & Davis A. Land Use, Land Cover, and Pollinator Health : A Review and Trend Analysis, Economic Research Report Number 232, 2017.
- Higuchi M, Honda T, Proske RJ, Yeh ET. Regulation of reactive oxygen species–induced apoptosis and necrosis by caspase 3–like proteases. *Oncogene*. 21 : 2753–2760, 1998.
- Ito K, Hirao A, Arai F, Takubo K, Matsuoka S, Miyamoto K, Ohmura M, Naka K, Hosokawa K, Ikeda Y & Suda T. Reactive oxygen species act through p38 MAPK to limit the lifespan of hematopoietic stem cells. *Nat Med* 12 : 446–451, 2006.
- Jaul E & Barron J. Age–Related Diseases and Clinical and Public Health Implications for the 85 Years Old and Over Population. *Front Public Health* 5 : 335, 2017.
- Jin TY, Saravanakumar K & Wang MH. In vitro and in vivo antioxidant properties of water and methanol extracts of linden bee pollen. *Biocatalysis and agricultural biotechnology* 13 : 186–189, 2018.
- Karabagias IK, Karabagias VK, Gatzias I & Riganakos KA. Bio–functional properties of bee pollen : The case of “bee pollen yoghurt”. *Coatings* 8 : 423, 2018.
- Kaškonienė V, Ruočkusienė G, Kaškonas P, Akuneca I & Maruška A. Chemometric analysis of bee pollen based on volatile and phenolic compound compositions and antioxidant properties. *Food Analytical Methods* 8 : 1150–1163, 2015.
- Ketkar SS, Rathore AS, Lohidasan S, Rao L, Paradkar AR & Mahadik KR. Investigation of the nutraceutical potential of monofloral Indian mustard bee pollen. *Journal of integrative medicine* 12 : 379–389, 2014.
- Khurelchuluun A, Uehara O, Paudel D, Morikawa T, Kawano Y, Sakata M, Shibata H, Yoshida K, Sato J, Mura H, Nagayasu H & Abiko Y. Bee Pollen Diet Alters the Bacterial Flora and Antimicrobial Peptides in the Oral Cavities of Mice. *Foods* 10 : 1282, 2021.
- Kolayli S, Ebru Cakir H & Sahin H. Anti–Inflammatory Activities of Some Bee Products by Inhibition of Bovine Testes Hyaluronidase. *Curr* 12 : 183–187, 2016.
- Kostić AŽ, Milinčić DD, Gašić UM, Nedić N, Stanojević SP, Tešić ŽL & Pešić MB. Polyphenolic profile and antioxidant properties of bee–collected pollen from sunflower (*Helianthus annuus* L.) plant. *LWT* 112 : 108244, 2019.
- Kunugi H & Mohammed AA. Royal Jelly and Its Components Promote Healthy Aging and Longevity : From Animal Models to Humans. *Int. J. Mol. Sci*. 20 : 4662, 2019.
- Kurek–Górecka A, Górecki M, Rzepecka–Stojko A, Balwierz R & Stojko J. Bee Products in Dermatology and Skin Care. *Molecules* 25 : 556, 2020.
- Lahiri D, Nag M, Banerjee R, Mukherjee D, Garai S, Sarkar T, Dey A, Sheikh HI, Pathak SK, Edinur HA, Pati S & Ray RR. Amylases : Biofilm Inducer or Biofilm Inhibitor? *Front. Cell. Infect. Microbiol*. 11 : 355, 2021.
- LeBlanc BW, Davis OK, Boue S, DeLucca A, Deeby T. Antioxidant activity of Sonoran Desert bee pollen. *Food chemistry* 115 : 1299–1305, 2009.
- Lee KS, Lee BS, Semnani S, Avanesian A, Um CY, Jeon HJ, Seong KM, Yu K, Min KJ & Jafari M. Curcumin Extends Life Span, Improves Health Span, and Modulates the Expression of Age–Associated Aging Genes in *Drosophila melanogaster*. *Rejuvenation Res*. 13 : 561–570, 2010.
- Leja M, Mareczek A, Wyżgolik G, Klepacz–Baniak J & Czekońska K. Antioxidative properties of bee pollen in selected plant species. *Food Chemistry* 100 : 237–240, 2007.
- Liu X & Li L. Morphological observation of effect of bee

- pollen on intercellular lipofuscin in NIH mice. *Zhongguo Zhongyao Zazhi* 15 : 561–578, 1990.
- Luceri C, Bigagli E, Pitozzi V & Giovannelli L. A nutrigenomics approach for the study of anti-aging interventions : olive oil phenols and the modulation of gene and microRNA expression profiles in mouse brain. *Eur J Nutr*. 56 : 865–877, 2017.
- Mandal SM, Dias RO & Franco OL. Phenolic Compounds in Antimicrobial Therapy. *J. Med. Food* 20 : 1031–1038, 2017.
- Middleton E & Kandaswami C. Effects of flavonoids on immune and inflammatory cell functions. *Biochem. Pharmacol.* 43 : 1167–1179, 1992.
- Middleton E. Effect of plant flavonoids on immune and inflammatory cell function. *Adv Exp Med Biol* 439 : 175–182, 1998.
- Mohdaly AA, Mahmoud AA, Roby MH, Smetanska I & Ramadan MF. Phenolic extract from propolis and bee pollen : composition, antioxidant and antibacterial activities. *Journal of Food Biochemistry* 39 : 538–547, 2015.
- Morais M, Moreira L, Feás X & Estevinho LM. Honeybee-collected pollen from five Portuguese Natural Parks : Palynological origin, phenolic content, antioxidant properties, and antimicrobial activity. *Food Chem. Toxicol.* 49 : 1096–1101, 2011.
- Morgia G, Falsaperla M, Malaponte G, Madonia M, Indelicato M, Travali S & Mazzarino MC. Matrix metalloproteinases as diagnostic (MMP-13) and prognostic (MMP-2, MMP-9) markers of prostate cancer. *Urol Res* 33 : 44–50, 2005.
- Nogueira C, Iglesias A, Feás X & Estevinho LM. Commercial Bee Pollen with Different Geographical Origins : A Comprehensive Approach. *Int. J. Mol. Sci.* 13 : 11173–11187, 2012.
- Nuvoloni R, Meucci V, Turchi B, Sagona S, Fratini F, Felicioli A, Cerri D & Pedonese F. Bee-pollen retailed in Tuscany (Italy) : labelling, palynological, microbiological, and mycotoxicological profile. *LWT – LWT-FOOD SCI TECHNOL*, 2020.
- Oliveira BF, Nogueira-Machado JA & Chaves MM. The Role of Oxidative Stress in the Aging Process. *ScientificWorldJournal* 10 : 1121–1128, 2010.
- Omar WA, Azhar NA, Harif Fadzilah N & Nik Mohamed Kamal NNS. Bee pollen extract of Malaysian stingless bee enhances the effect of cisplatin on breast cancer cell lines. *Asian Pac J Cancer Prev* 6 : 265–269, 2016.
- Pasupuleti VR, Sammugam L, Ramesh N & Gan SH. Honey, Propolis, and Royal Jelly : A Comprehensive Review of Their Biological Actions and Health Benefits. *Oxid Med Cell Longev* e1259510, 2017.
- Peng C, Zuo Y, Kwan KM, Liang Y, Ma KY, Chan HYE, Huang Y, Yu H & Chen ZY. Blueberry extract prolongs lifespan of *Drosophila melanogaster*. *Exp. Gerontol.* 47 : 170–178, 2012.
- Petri S, Körner S & Kiaei M. Nrf2/ARE Signaling Pathway : Key Mediator in Oxidative Stress and Potential Therapeutic Target in ALS. *Neurol. Res. Int.* e878030, 2012.
- Pham CG, Bubici C, Zazzeroni F, Papa S, Jones J, Alvarez K, Jayawardena S, De Smaele E, Cong R, Beaumont C, Torti FM, Torti SV & Franzoso G. Ferritin Heavy Chain Upregulation by NF- κ B Inhibits TNF α -Induced Apoptosis by Suppressing Reactive Oxygen Species. *Cell* 119 : 529–542, 2004.
- Pietsch K, Saul N, Swain SC, Menzel R, Steinberg CW & Stürzenbaum SR. Meta-Analysis of Global Transcriptomics Suggests that Conserved Genetic Pathways are Responsible for Quercetin and Tannic Acid Mediated Longevity in *C. elegans*. *Front Genet* 3 : 48, 2012.
- Rescigno T, Micolucci L, Tecce MF & Capasso A. Bioactive Nutrients and Nutrigenomics in Age-Related Diseases. *Molecules* 22 : 105, 2017.
- Rocchetti G, Castiglioni S, Maldarizzi G, Carloni P & Lucini L. UHPLC-ESI-QTOF-MS phenolic profiling and antioxidant capacity of bee pollen from different botanical origin. *International Journal of Food Science & Technology* 54 : 335–346, 2019.
- Rzepecka-Stojko A, Stojko J & Kurek-Górecka A. Polyphenols from Bee Pollen : Structure, Absorption, Metabolism and Biological Activity. *Molecules*. 20 : 21732–21749, 2015.
- Safar H, Van WJ, Møller P & Jacobsen C. Carotenoids, Phenolic Compounds and Tocopherols Contribute to the Antioxidative Properties of Some Microalgae Species Grown on Industrial Wastewater. *Mar Drugs* 13 : 7339–7356, 2015.
- Sajadimajd S & Khazaei M. Oxidative Stress and Cancer : The Role of Nrf2. *Curr Cancer Drug Targets* 18 : 538–557, 2018
- Singh R, Wang Y, Schattenberg JM, Xiang Y & Czaja MJ.

- Chronic oxidative stress sensitizes hepatocytes to death from 4-hydroxynonenal by JNK/c-Jun overactivation. *Am J Physiol Gastrointest Liver Physiol* 297 : G907–917, 2009.
- Soares de Arruda VA, Viera dos Santos A, Figueiredo Sampaio D, da Silva Araújo E, de Castro Peixoto AL, Estevinho LM & de Almeida–Muradian LB. Brazilian bee pollen : phenolic content, antioxidant properties and antimicrobial activity. *J. Apic. Res.* 60 : 775–783, 2021.
- Sun L, Guo Y, Zhang Y & Zhuang Y. Antioxidant and anti-tyrosinase activities of phenolic extracts from rape bee pollen and inhibitory melanogenesis by cAMP/MITF/TYR pathway in B16 mouse melanoma cells. *Frontiers in pharmacology* 8 : 104, 2017.
- Tanigawa S, Fujii M & Hou DX. Action of Nrf2 and Keap1 in ARE-mediated NQO1 expression by quercetin. *Free Radic. Biol. Med.* 42 : 1690–1703, 2007.
- Tomás–Lorente F, García–Grau MM, Nieto JL & Tomás–Barberán FA. Flavonoids from *Cistus ladanifer* bee pollen. *Phytochemistry* 31 : 2027–2029, 1992.
- Uçar M, Değir O, Gerigelmez AY, Cengiz S, Barlak Y & Ovalı E. Effect of Turkish pollen and propolis extracts on caspase-3 activity in myeloid cancer cell lines. *Trop. J. Pharm. Res.* 15 : 2445–2449, 2016.
- Ulusoy E, Kolaylı S. Phenolic composition and antioxidant properties of Anzer bee pollen. *Journal of Food Biochemistry* 38 : 73–82, 2014.
- Vafadar A, Shabaninejad Z, Movahedpour A & Fallahi F. Quercetin and cancer : new insights into its therapeutic effects on ovarian cancer cells. *Cell Biosci* 10 : 10–32, 2020.
- Vasconcelos MR, Duarte AW, Gomes EP, Silva SC & López AM. Physicochemical composition and antioxidant potential of bee pollen from different botanical sources in Alagoas, Brazil. *Ciência e Agrotecnologia* 41 : 447–458, 2017.
- Weng CJ & Yen GC. Chemopreventive effects of dietary phytochemicals against cancer invasion and metastasis : Phenolic acids, monophenol, polyphenol, and their derivatives. *Cancer Treat. Rev.* 38 : 76–87, 2012.
- Wojdyło A, Nowicka P & Bąbalewski P. Phenolic and carotenoid profile of new goji cultivars and their anti-hyperglycemic, anti-aging and antioxidant properties. *J. Funct. Foods* 48 : 632–642, 2018.
- Xia X, Xiang X, Huang F, Zheng M, Cong R, Han L & Zhang Z. Dietary polyphenol canolol from rapeseed oil attenuates oxidative stress-induced cell damage through the modulation of the p38 signaling pathway. *RSC Advances* 8 : 24338–24345, 2018.
- Yahfoufi N, Alsadi N, Jambi M & Matar C. The Immunomodulatory and Anti-Inflammatory Role of Polyphenols. *Nutrients* 10 : E1618, 2018.
- Yasumoto R, Kawanishi H, Tsujino T, Tsujita M, Nishisaka N, Horii A & Kishimoto T. Clinical evaluation of long-term treatment using cernitin pollen extract in patients with benign prostatic hyperplasia. *Clin. Ther.* 17 : 82–87, 1995.
- Yue Y, Shen P, Xu Y & Park Y. p-Coumaric acid improves oxidative and osmotic stress responses in *Caenorhabditis elegans*. *J. Sci. Food Agric.* 99 : 1190–1197, 2019.
- Zhang Y, Yang F, Jamali MA & Peng Z. Antioxidant enzyme activities and lipid oxidation in rape (*Brassica campestris* L.) bee pollen added to salami during processing. *Molecules* 21 : 1439, 2016.
- Zuluaga C, Martínez A, Fernández J, López–Baldó J, Quiles A & Rodrigo D. Effect of high pressure processing on carotenoid and phenolic compounds, antioxidant capacity, and microbial counts of bee-pollen paste and bee-pollen-based beverage. *Innovative Food Science & Emerging Technologies* 37 : 10–17, 2016.